

DermaPlast® EFFECT XL, Heel & Small Blisters

General Product Description

DermaPlast EFFECT XL Blisters presentations are composed of a layer of 600 µm / 0,6 mm hydrocolloid material acting as the wound pad, another larger layer of 100 µm / 0,1 mm hydrocolloid material acting as the dressing backing material, the silicone paper protection in contact with the adhesive coated side of the dressing and the peel pack (individually sealed plaster).

DermaPlast EFFECT Heel Blisters and DermaPlast EFFECT Small Blisters presentations are composed of a layer of 300 µm / 0,3 mm hydrocolloid material acting as the wound pad, another larger layer of 300 µm / 0,3 mm hydrocolloid material acting as the dressing backing material, the silicone paper protection in contact with the adhesive coated side of the dressing and the peel pack (individually sealed plaster).

All individually sealed plasters (one plaster in each peel pack) are packed in a labeled plastic folding box, which includes a leaflet with instructions for use.

Intended Purpose

Hydrocolloid plasters are non-active, single-use, self-adhesive, non-sterile devices consisting of a self-adhesive hydrocolloid wound contact layer and an adhesive coated backing material (carrier material), protected with a silicone paper, for the protection and treatment of blisters.

The products are intended for humans, without restrictions (in age group, weight range, health status or condition), for use by lay-users, in contact with intact skin (healthy skin) and breached skin, and short-term (60 minutes ≤ Duration ≤ 30 days, acc. to MDR) / prolonged use (24 hours < Duration ≤ 30 days, acc. to ISO 10993-1:2018) duration of use.

The products do not contain resorbable components, medicinal products / drug substances (derived from human blood or plasma or not), tissues or cells of human or animal origin, radioactive substances, nanomaterials or other hazardous components.

Application/Indication

- Wash hands and ensure the affected area is clean, dry and oil free before applying the plaster.
- Open the peel pack and extract the plaster.
- Warm the plaster with the hands before applying.
- Remove the first protective paper and apply the exposed part of the plaster to the affected skin area.
- Then gently pull the second protective paper and apply the rest of the plaster. Gently mould the plaster edges to ensure optimum adhesion.
- The application maximum of the hydrocolloid is up to 4 days per individual product. Should you wish to remove the plaster early, first wet it briefly with warm water.

DermaPlast® EFFECT XL, Heel & Small Blisters

Reference Numbers

Ref no.	Description	Length	Content of folding box
5362401	DermaPlast EFFECT XL Blister plaster	75x45mm	P6
5361401	DermaPlast EFFECT Heel	76x45mm	P6
5363401	DermaPlast EFFECT Blister small	55x19mm	P6

Residual Risks, Contra-Indications and Undesirable Side Effects, Warnings

The hydrocolloid plasters must not be used on non-treated infected blisters, highly bleeding or exudative wounds or for pressure ulcers.

Adverse reactions may occur in patients who are allergic or hypersensitive to the product components.

All presentations:

- Only apply the plaster if the hydrocolloid pad completely covers the blister/minor injury/corn/cold sore/abrasion.
- In cases of diabetes mellitus always consult a doctor before using plasters.
- Do not apply the product for more than 4 days.
- Store plasters in a cool (not in the fridge), dry place, out of the reach of children.

All presentations except Cold sore presentations:

- Consult a doctor before applying the plaster on a potentially infected blister, minor injury, or if the blister or minor injury shows signs of infection while plasters are in place (e.g. redness, overheating, swelling or pain).

Specially for Blister presentations:

- Do not burst a sealed blister as this could risk an infection.

Single Use Device

Reusing a disposable and single-use medical device is dangerous and it could cause risk of infection. Reprocessing devices, in order to reuse them, may seriously damage their integrity and their performance. Information available on request.

Product Disposal

To minimize the risk of potential infection hazards, or environmental pollution, disposable components of DermaPlast® EFFECT XL, Heel & Small Blisters should follow disposal procedures according to applicable and local laws, rules, regulations and infection prevention standards.

DermaPlast® EFFECT XL, Heel & Small Blisters

Incident Reporting

For a patient/user/third party in the European Union and in countries with identical regulatory regime (Regulation 2017/745/EU on Medical Devices); if, during the use of this device or as a result of its use, a serious incident has occurred, please report it to the manufacturer and/or its authorised representative and to your national authority.

Product Performance /Material Characteristics

Formulation Hydrocolloids 300 µm/0,3 mm			
Property	Test method	Unit	Requirements
Peel adhesion on Steal Plates (90°)	TM00 0090-04	N/25mm	6,9 (+2,7 / - 4,1)

Formulation Hydrocolloids 600 µm/0,6 mm			
Property	Test method	Unit	Requirements
Peel adhesion on Steal Plates (90°)	TM00 0090-04	N/25mm	8,4 (+2,7 / - 4,9)

Labelling

Lot-No. with 9-Digit Code

e.g.:



0XXX

XX

12X

Year and number of production

Week of production

For internal purposes only

Manufacturing Date

e.g.:



2020
year

01
month

07
day

Use-by-Date

e.g.:



2023 - 01
year month

Shelf Life: 3 years

Medical Device



DermaPlast® EFFECT XL, Heel & Small Blisters

Unique Device Identification (UDI)



Distributor



Latest Date of Revision: 2020-03-12